

ELLIPTIC CURVES OVER THE PERFECT CLOSURE OF A FUNCTION FIELD

DRAGOS GHIOCA

ABSTRACT. We prove that the group of rational points of a non-isotrivial elliptic curve defined over the perfect closure of a function field in one variable over a finite field is finitely generated.

1. INTRODUCTION

The main ingredient for our result is a study of the Lehmer inequality for elliptic curves. The classical Lehmer conjecture (see [7], page 476) asserts that there is an absolute constant $C > 0$ so that any algebraic number α that is not a root of unity satisfies the following inequality for its logarithmic height

$$h(\alpha) \geq \frac{C}{[\mathbb{Q}(\alpha) : \mathbb{Q}]}.$$

A partial result towards this conjecture is obtained in [3]. The analog of Lehmer conjecture for elliptic curves and abelian varieties asks for a good lower bound for the canonical height of a non-torsion point of the abelian variety. Also this question has been much studied (see [1], [2], [5], [9], [15]). As a consequence of our work, we obtain the following result.

Theorem 1.1. *Let K be a function field of transcendence degree 1 over $\mathbb{F}_p$. Let E be a non-isotrivial elliptic curve defined over K . Then $E(K^{\text{per}})$ is finitely generated.*

Using completely different methods, Minhyong Kim studied the set of rational points of non-isotrivial curves of genus at least two over the perfect closure of a function field in one variable over a finite field (see [6]).

Using the result of Theorem 1.1, Thomas Scanlon proved the full positive characteristic Mordell-Lang Conjecture for abelian varieties that are isogenous to a product of elliptic curves (see [11]).

Our plan is to prove first some general results about tame modules in Section 2 and then to prove Theorem 1.1 in Section 3.

2. TAME MODULES

Definition 2.1. Let R be an integral domain and let K be its field of fractions. If M is an R -module, then by the *rank* of M , denoted $\text{rk}(M)$, we mean the dimension of the K -vector space $M \otimes_R K$. We call M a *tame* module if every finite rank submodule of M is finitely generated.

Lemma 2.2. *Let R be a Dedekind domain and let M be an R -module. Assume there exists a function $h : M \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following properties*

- (i) (triangle inequality) $h(x \pm y) \leq h(x) + h(y)$, for every $x, y \in M$.

- (ii) if $x \in M_{\text{tor}}$, then $h(x) = 0$.
 - (iii) there exists $c > 0$ such that for each $x \notin M_{\text{tor}}$, $h(x) > c$.
 - (iv) there exists $a \in R \setminus \{0\}$ such that R/aR is finite and for all $x \in M$, $h(ax) \geq 4h(x)$.
- If M_{tor} is finite, then M is a tame R -module.

Proof. By the definition of a tame module, it suffices to assume that M is a finite rank R -module and conclude that it is finitely generated.

Let $a \in R$ as in (iv) of Lemma 2.2. By Lemma 3 of [10], M/aM is finite (here we use the assumption that M_{tor} is finite). The following result is the key to the proof of Lemma 2.2.

Sublemma 2.3. For every $D > 0$, there exists finitely many $x \in M$ such that $h(x) \leq D$.

Proof of Sublemma 2.3. If we suppose Sublemma 2.3 is not true, then we can define

$$C = \inf\{D \mid \text{there exists infinitely many } x \in M \text{ such that } h(x) \leq D\}.$$

Properties (ii) and (iii) and the finiteness of M_{tor} yield $C \geq c > 0$. By the definition of C , it must be that there exists an infinite sequence of elements z_n of M such that for every n ,

$$h(z_n) < \frac{3C}{2}.$$

Because M/aM is finite, there exists a coset of aM in M containing infinitely many z_n from the above sequence.

But if $k_1 \neq k_2$ and z_{k_1} and z_{k_2} are in the same coset of aM in M , then let $y \in M$ be such that $ay = z_{k_1} - z_{k_2}$. Using properties (iv) and (i), we get

$$h(y) \leq \frac{h(z_{k_1} - z_{k_2})}{4} \leq \frac{h(z_{k_1}) + h(z_{k_2})}{4} < \frac{3C}{4}.$$

We can do this for any two elements of the sequence that lie in the same coset of aM in M . Because there are infinitely many of them lying in the same coset, we can construct infinitely many elements $z \in M$ such that $h(z) < \frac{3C}{4}$, contradicting the minimality of C . $\square$

From this point on, our proof of Lemma 2.2 follows the classical descent argument in the Mordell-Weil theorem (see [12]).

Take coset representatives $y_1, \dots, y_k$ for aM in M . Define then

$$B = \max_{i \in \{1, \dots, k\}} h(y_i).$$

Consider the set $Z = \{x \in M \mid h(x) \leq B\}$, which is finite according to Sublemma 2.3. Let N be the finitely generated R -submodule of M which is spanned by Z .

We claim that $M = N$. If we suppose this is not the case, then by Sublemma 2.3 we can pick $y \in M - N$ which minimizes $h(y)$. Because N contains all the coset representatives of aM in M , we can find $i \in \{1, \dots, k\}$ such that $y - y_i \in aM$. Let $x \in M$ be such that $y - y_i = ax$. Then $x \notin N$ because otherwise it would follow that $y \in N$ (we already know $y_i \in N$). By our choice of y and by properties (iv) and (i), we have

$$h(y) \leq h(x) \leq \frac{h(y - y_i)}{4} \leq \frac{h(y) + h(y_i)}{4} \leq \frac{h(y) + B}{4}.$$

This means that $h(y) \leq \frac{B}{3} < B$. This contradicts the fact that $y \notin N$ because N contains all the elements $z \in M$ such that $h(z) \leq B$. This contradiction shows that indeed $M = N$ and so, M is finitely generated. $\square$

Corollary 2.4. *Let R be a Dedekind domain and let M be a tame R -module.*

(a) *If $\text{rk}(M) = \aleph_0$, then M is a direct sum of a finite torsion submodule and a free submodule of rank $\aleph_0$.*

(b) *If $\text{rk}(M)$ is finite and R is a principal ideal domain, then M is a direct sum of a finite torsion submodule and a free submodule of finite rank.*

Proof. Part (a) of Corollary 2.4 is proved in Proposition 10 of [10].

If $\text{rk}(M)$ is finite and because M is tame, we conclude that M is finitely generated. Because R is a principal ideal domain we get the result of part (b) of Corollary 2.4. $\square$

3. ELLIPTIC CURVES

The setting is the following: K is a finitely generated field of transcendence degree 1 over $\mathbb{F}_p$ where p is a prime as always. We fix an algebraic closure K^{alg} of K . We denote by $\mathbb{F}_p^{\text{alg}}$ the algebraic closure of $\mathbb{F}_p$ inside K^{alg} .

Let E be a non-isotrivial elliptic curve (i.e. $j(E) \notin \mathbb{F}_p^{\text{alg}}$) defined over K . Let K^{per} be the perfect closure of K inside K^{alg} . We will prove in Theorem 1.1 that $E(K^{\text{per}})$ is finitely generated.

For every finite extension L of K we denote by M_L the set of discrete valuations v on L , normalized so that the value group of v is $\mathbb{Z}$. For each $v \in M_L$ we denote by f_v the degree of the residue field of v over $\mathbb{F}_p$. If $P \in E(L)$ and $m \in \mathbb{Z}$, mP represents the point on the elliptic curve obtained using the group law on E . We define a notion of height for the point $P \in E(L)$ with respect to the field K (see Chapter *VIII* of [13] and Chapter *III* of [14])

$$(1) \quad h_K(P) = \frac{-1}{[L : K]} \sum_{v \in M_L} f_v \min\{0, v(x(P))\}.$$

Then we define the canonical height of P with respect to K as

$$(2) \quad \hat{h}_{E/K}(P) = \frac{1}{2} \lim_{n \rightarrow \infty} \frac{h_K(2^n P)}{4^n}.$$

We also denote by $\Delta_{E/K}$ the divisor which is the minimal discriminant of E with respect to the field K (see Chapter *VIII* of [13]). By $\deg(\Delta_{E/K})$ we denote the degree of the divisor $\Delta_{E/K}$ (computed with respect to $\mathbb{F}_p$). We denote by g_K the genus of the function field K .

The following result is proved in [4].

Theorem 3.1 (Goldfeld-Szpiro). *Let E be an elliptic curve over a function field K of one variable over a field in any characteristic. Let $\hat{h}_{E/K}$ denote the canonical height on E and let $\Delta_{E/K}$ be the minimal discriminant of E , both computed with respect to K . Then for every point $P \in E(K)$ which is not a torsion point:*

$$\hat{h}_{E/K}(P) \geq 10^{-13} \deg(\Delta_{E/K}) \text{ if } \deg(\Delta_{E/K}) \geq 24(g_K - 1),$$

and

$$\hat{h}_{E/K}(P) \geq 10^{-13-23g} \deg(\Delta_{E/K}) \text{ if } \deg(\Delta_{E/K}) < 24(g_K - 1).$$

We are ready to prove our result.

Proof of Theorem 1.1. We first observe that replacing K by a finite extension does not affect the conclusion of the theorem. Thus, at the expense of replacing K by a finite extension, we

may assume E is semi-stable over K (the existence of such a finite extension is guaranteed by Proposition 5.4 (a) of [13]).

As before, we let $\hat{h}_{E/K}$ and $\Delta_{E/K}$ be the canonical height on E and the minimal discriminant of E , respectively, computed with respect to K .

For every $n \geq 1$, we denote by $E^{(p^n)}$ the elliptic curve $F^n(E)$, where F is the usual Frobenius (seen as morphism of varieties). Thus

$$(3) \quad F^n : E(K^{1/p^n}) \rightarrow E^{(p^n)}(K)$$

is a bijection. Moreover, for every $P \in E(K^{1/p})$,

$$(4) \quad pP = (VF)(P) \in V(E^{(p)}(K)) \subset E(K)$$

where V is the Verschiebung. Similarly, we get that

$$(5) \quad p^n E(K^{1/p^n}) \subset E(K) \text{ for every } n \geq 1.$$

Thus $E(K^{\text{per}})$ lies in the p -division hull of the $\mathbb{Z}$ -module $E(K)$. Because $E(K)$ is finitely generated (by the Mordell-Weil theorem), we conclude that $E(K^{\text{per}})$, as a $\mathbb{Z}$ -module, has finite rank.

We will show next that the height function $\hat{h}_{E/K}$ and $p \in \mathbb{Z}$ satisfy the properties (i)-(iv) of Lemma 2.2. Properties (i) and (ii) are well-known for $\hat{h}_{E/K}$ and we also have the formula (see Chapter VIII of [13])

$$\hat{h}_{E/K}(pP) = p^2 \hat{h}_{E/K}(P) \geq 4 \hat{h}(P) \text{ for every } P \in E(K^{\text{alg}}),$$

which proves that property (iv) of Lemma 2.2 holds.

Let P be a non-torsion point of $E(K^{\text{per}})$. Then $P \in E(K^{1/p^n})$ for some $n \geq 0$. Because K^{1/p^n} is isomorphic to K , they have the same genus, which we call it g . We denote by $\hat{h}_{E/K^{1/p^n}}$ and $\Delta_{E/K^{1/p^n}}$ the canonical height on E and the minimal discriminant of E , respectively, computed with respect to K^{1/p^n} . Using Theorem 3.1, we conclude

$$(6) \quad \hat{h}_{E/K^{1/p^n}}(P) \geq 10^{-13-23g} \deg(\Delta_{E/K^{1/p^n}}).$$

We have $\hat{h}_{E/K^{1/p^n}}(P) = [K^{1/p^n} : K] \hat{h}_{E/K}(P) = p^n \hat{h}_{E/K}(P)$. Similarly, using the proof of Proposition 5.4 (b) of [13],

$$\deg(\Delta_{E/K^{1/p^n}}) = p^n \deg(\Delta_{E/K}).$$

We conclude that for every $P \in E(K^{\text{per}})$,

$$(7) \quad \hat{h}_{E/K}(P) \geq 10^{-13-23g} \deg(\Delta_{E/K}).$$

Because E is non-isotrivial, $\Delta_{E/K} \neq 0$ and so, $\deg(\Delta_{E/K}) \geq 1$. We conclude

$$(8) \quad \hat{h}_{E/K}(P) \geq 10^{-13-23g}.$$

Inequality (8) shows that property (iii) of Lemma 2.2 holds for $\hat{h}_{E/K}$. Thus properties (i)-(iv) of Lemma 2.2 hold for $\hat{h}_{E/K}$ and $p \in \mathbb{Z}$.

We show that $E_{\text{tor}}(K^{\text{per}})$ is finite. Equation (5) shows that the prime-to- p -torsion of $E(K^{\text{per}})$ equals the prime-to- p -torsion of $E(K)$; thus the prime-to- p -torsion of $E(K^{\text{per}})$ is finite. If there exists infinite p -power torsion in $E(K^{\text{per}})$, equation (3) yields that we have arbitrarily large p -power torsion in the family of elliptic curves $E^{(p^n)}$ over K . But this contradicts standard results on uniform boundedness for the torsion of elliptic curves over

function fields, as established in [8] (actually, [8] proves a uniform boundedness of the entire torsion of elliptic curves over a fixed function field; thus including the prime-to- p -torsion). Hence $E_{\text{tor}}(K^{\text{per}})$ is finite.

Because all the hypothesis of Lemma 2.2 hold, we conclude that $E(K^{\text{per}})$ is tame. Because $\text{rk}(E(K^{\text{per}}))$ is finite we conclude by Corollary 2.4 (b) that $E(K^{\text{per}})$ is a direct sum of a finite torsion submodule and a free submodule of finite rank. $\square$

Remark 3.2. It is absolutely crucial in Theorem 1.1 that E is non-isotrivial. Theorem 1.1 fails in the isotrivial case, i.e. there exists no $n \geq 0$ such that $E(K^{\text{per}}) = E(K^{1/p^n})$. Indeed, if E is defined by $y^2 = x^3 + x$ ($p > 2$), $K = \mathbb{F}_p(t, (t^3 + t)^{\frac{1}{2}})$ and $P = (t, (t^3 + t)^{\frac{1}{2}})$, then $F^{-n}P \in E(K^{1/p^n}) \setminus E(K^{1/p^{n-1}})$, for every $n \geq 1$. So, $E(K^{\text{per}})$ is not finitely generated in this case (and we can get a similar example also for the case $p = 2$).

REFERENCES

- [1] M. Baker, J. Silverman, *A lower bound for the canonical height on abelian varieties over abelian extensions*. Math. Res. Lett. 11 (2004), no. 2-3, 377-396.
- [2] S. David, J. Silverman, *Minoration de la hauteur de Néron-Tate sur les variétés abéliennes de type C*. M. (French) [Lower bound for the Néron-Tate height on abelian varieties of CM type] J. Reine Angew. Math. 529 (2000), 1-74.
- [3] E. Dobrowolski, *On a question of Lehmer and the number of irreducible factors of a polynomial*. Acta Arith. 34, no. 4, 391-401, (1979).
- [4] D. Goldfeld, L. Szpiro, *Bounds for the order of the Tate-Shafarevich group* Special issue in honour of Frans Oort. Compositio Math. 97 (1995), no. 1-2, 71-87
- [5] M. Hindry, J. Silverman, *On Lehmer's conjecture for elliptic curves*. Séminaire de Théorie des Nombres, Paris 1988-1989, 103-116, Progr. Math., No. 91, Birkhäuser Boston, Boston, MA, 1990.
- [6] Minhyong Kim, *Purely inseparable points on curves of higher genus*. Math.Res.Lett. 4 (1997), no. 5, 663-666.
- [7] D. H. Lehmer, *Factorization of certain cyclotomic polynomials*, Ann. of Math. (2) 34 (1933), no. 3, 461-479.
- [8] Martin Levin, *On the group of rational points on elliptic curves over function fields*, Amer.J.Math 90, 1968 ,456-462
- [9] D.W. Masser, *Counting points of small height on elliptic curves*. Bull. Soc. Math. France 117 (1989), no. 2, 247-265.
- [10] B. Poonen, *Local height functions and the Mordell-Weil theorem for Drinfeld modules*, Compositio Mathematica **97** (1995), 349-368.
- [11] T. Scanlon, *Positive characteristic Manin-Mumford theorem*. preprint, 2003, available online at <http://Math.Berkeley.EDU/~scanlon/papers/papers.html>.
- [12] Jean-Pierre Serre, *Lectures on the Mordell-Weil theorem*. Translated from the French and edited by Martin Brown from notes by Michel Waldschmidt. Aspects of Mathematics, E15. Friedr. Vieweg & Sohn, Braunschweig, 1989. x+218 pp.
- [13] Joseph Silverman, *The arithmetic of elliptic curves*. Graduate Texts in Mathematics, 106. Springer-Verlag, New York, 1986. xii+400 pp.
- [14] Joseph Silverman, *Advanced topics in the arithmetic of elliptic curves*. Graduate Texts in Mathematics, 151. Springer-Verlag, New York, 1994. xiv+525 pp.
- [15] J. Silverman, *A lower bound for the canonical height on elliptic curves over abelian extensions*. J. Number Theory 104 (2004), no. 2, 353-372.